

### 3. Double Integrals

#### 3A. Double integrals in rectangular coordinates

##### 3A-1

a) Inner:  $6x^2y + y^2 \Big|_{y=-1}^1 = 12x^2$ ; Outer:  $4x^3 \Big|_0^2 = 32$ .

b) Inner:  $-u \cos t + \frac{1}{2}t^2 \cos u \Big|_{t=0}^{\pi} = 2u + \frac{1}{2}\pi^2 \cos u$   
 Outer:  $u^2 + \frac{1}{2}\pi^2 \sin u \Big|_0^{\pi/2} = (\frac{1}{2}\pi)^2 + \frac{1}{2}\pi^2 = \frac{3}{4}\pi^2$ .

c) Inner:  $x^2y^2 \Big|_{\sqrt{x}}^{x^2} = x^6 - x^3$ ; Outer:  $\frac{1}{7}x^7 - \frac{1}{4}x^4 \Big|_0^1 = \frac{1}{7} - \frac{1}{4} = -\frac{3}{28}$

d) Inner:  $v\sqrt{u^2+4} \Big|_0^u = u\sqrt{u^2+4}$ ; Outer:  $\frac{1}{3}(u^2+4)^{3/2} \Big|_0^1 = \frac{1}{3}(5\sqrt{5}-8)$

##### 3A-2

a) (i)  $\iint_R dy dx = \int_{-2}^0 \int_{-x}^2 dy dx$  (ii)  $\iint_R dx dy = \int_0^2 \int_{-y}^0 dx dy$

b) i) The ends of  $R$  are at 0 and 2, since  $2x - x^2 = 0$  has 0 and 2 as roots.



$$\iint_R dy dx = \int_0^2 \int_0^{2x-x^2} dy dx$$

ii) We solve  $y = 2x - x^2$  for  $x$  in terms of  $y$ : write the equation as  $x^2 - 2x + y = 0$  and solve for  $x$  by the quadratic formula, getting  $x = 1 \pm \sqrt{1-y}$ . Note also that the maximum point of the graph is (1, 1) (it lies midway between the two roots 0 and 2). We get

$$\iint_R dx dy = \int_0^1 \int_{1-\sqrt{1-y}}^{1+\sqrt{1-y}} dx dy,$$

c) (i)  $\iint_R dy dx = \int_0^{\sqrt{2}} \int_0^x dy dx + \int_{\sqrt{2}}^2 \int_0^{\sqrt{4-x^2}} dy dx$

(ii)  $\iint_R dx dy = \int_0^{\sqrt{2}} \int_y^{\sqrt{4-y^2}} dx dy$



d) Hint: First you have to find the points where the two curves intersect, by solving simultaneously  $y^2 = x$  and  $y = x - 2$  (eliminate  $x$ ).

The integral  $\iint_R dy dx$  requires two pieces;  $\iint_R dx dy$  only one.

3A-3 a)  $\iint_R x dA = \int_0^2 \int_0^{1-x/2} x dy dx$ ;

Inner:  $x(1 - \frac{1}{2}x)$  Outer:  $\frac{1}{2}x^2 - \frac{1}{6}x^3 \Big|_0^2 = \frac{4}{2} - \frac{8}{6} = \frac{2}{3}$ .

$$\text{b) } \iint_R (2x + y^2) dA = \int_0^1 \int_0^{1-y^2} (2x + y^2) dx dy$$

Inner:  $x^2 + y^2 x \Big|_0^{1-y^2} = 1 - y^2$ ;      Outer:  $y - \frac{1}{3}y^3 \Big|_0^1 = \frac{2}{3}$ .

$$\text{c) } \iint_R y dA = \int_0^1 \int_{y-1}^{1-y} y dx dy$$

Inner:  $xy \Big|_{y-1}^{1-y} = y[(1-y) - (y-1)] = 2y - 2y^2$       Outer:  $y^2 - \frac{2}{3}y^3 \Big|_0^1 = \frac{1}{3}$ .

$$\text{3A-4 a) } \iint_R \sin^2 x dA = \int_{-\pi/2}^{\pi/2} \int_0^{\cos x} \sin^2 x dy dx$$

Inner:  $y \sin^2 x \Big|_0^{\cos x} = \cos x \sin^2 x$       Outer:  $\frac{1}{3} \sin^3 x \Big|_{-\pi/2}^{\pi/2} = \frac{1}{3}(1 - (-1)) = \frac{2}{3}$ .



$$\text{b) } \iint_R xy dA = \int_0^1 \int_{x^2}^x (xy) dy dx.$$

Inner:  $\frac{1}{2}xy^2 \Big|_{x^2}^x = \frac{1}{2}(x^3 - x^5)$       Outer:  $\frac{1}{2} \left( \frac{x^4}{4} - \frac{x^6}{6} \right) \Big|_0^1 = \frac{1}{2} \cdot \frac{1}{12} = \frac{1}{24}$ .

c) The function  $x^2 - y^2$  is zero on the lines  $y = x$  and  $y = -x$ , and positive on the region  $R$  shown, lying between  $x = 0$  and  $x = 1$ . Therefore

$$\text{Volume} = \iint_R (x^2 - y^2) dA = \int_0^1 \int_{-x}^x (x^2 - y^2) dy dx.$$

$$\text{Inner: } x^2y - \frac{1}{3}y^3 \Big|_{-x}^x = \frac{4}{3}x^3; \quad \text{Outer: } \frac{1}{3}x^4 \Big|_0^1 = \frac{1}{3}.$$



$$\text{3A-5 a) } \int_0^2 \int_x^2 e^{-y^2} dy dx = \int_0^2 \int_0^y e^{-y^2} dx dy = \int_0^2 e^{-y^2} y dy = -\frac{1}{2}e^{-y^2} \Big|_0^2 = \frac{1}{2}(1 - e^{-4})$$

$$\text{b) } \int_0^{\frac{1}{4}} \int_{\sqrt{t}}^{\frac{1}{2}} \frac{e^u}{u} du dt = \int_0^{\frac{1}{2}} \int_0^{u^2} \frac{e^u}{u} dt du = \int_0^{\frac{1}{2}} u e^u du = (u-1)e^u \Big|_0^{\frac{1}{2}} = 1 - \frac{1}{2}\sqrt{e}$$

$$\text{c) } \int_0^1 \int_{x^{1/3}}^1 \frac{1}{1+u^4} du dx = \int_0^1 \int_0^{u^3} \frac{1}{1+u^4} dx du = \int_0^1 \frac{u^3}{1+u^4} du = \frac{1}{4} \ln(1+u^4) \Big|_0^1 = \frac{\ln 2}{4}.$$

$$\text{3A-6 } 0; \quad 2 \iint_S e^x dA, \quad S = \text{right half of } R; \quad 4 \iint_Q x^2 dA, \quad Q = \text{first quadrant}$$

$$0; \quad 4 \iint_Q x^2 dA; \quad 0$$

$$\text{3A-7 a) } x^4 + y^4 \geq 0 \Rightarrow \frac{1}{1+x^4+y^4} \leq 1$$

$$\text{b) } \iint_R \frac{x dA}{1+x^2+y^2} \leq \int_0^1 \int_0^1 \frac{x}{1+x^2} dx dy = \frac{1}{2} \ln(1+x^2) \Big|_0^1 = \frac{\ln 2}{2} < \frac{.7}{2}.$$

### 3B. Double Integrals in polar coordinates

#### 3B-1

a) In polar coordinates, the line  $x = -1$  becomes  $r \cos \theta = -1$ , or  $r = -\sec \theta$ . We also need the polar angle of the intersection points; since the right triangle is a 30-60-90 triangle (it has one leg 1 and hypotenuse 2), the limits are (no integrand is given):





$$\iint_R dr d\theta = \int_{2\pi/3}^{4\pi/3} \int_{-\sec \theta}^2 dr d\theta.$$

c) We need the polar angle of the intersection points. To find it, we solve the two equations  $r = \frac{3}{2}$  and  $r = 1 - \cos \theta$  simultaneously. Eliminating  $r$ , we get  $\frac{3}{2} = 1 - \cos \theta$ , from which  $\theta = 2\pi/3$  and  $4\pi/3$ . Thus the limits are (no integrand is given):

$$\iint_R dr d\theta = \int_{2\pi/3}^{4\pi/3} \int_{3/2}^{1-\cos \theta} dr d\theta.$$

d) The circle has polar equation  $r = 2a \cos \theta$ . The line  $y = a$  has polar equation  $r \sin \theta = a$ , or  $r = a \csc \theta$ . Thus the limits are (no integrand is given):

$$\iint_R dr d\theta = \int_{\pi/4}^{\pi/2} \int_{2a \cos \theta}^{a \csc \theta} dr d\theta.$$



$$\mathbf{3B-2} \text{ a) } \int_0^{\pi/2} \int_0^{\sin 2\theta} \frac{r dr d\theta}{r} = \int_0^{\pi/2} \sin 2\theta d\theta = -\frac{1}{2} \cos 2\theta \Big|_0^{\pi/2} = -\frac{1}{2}(-1 - 1) = 1.$$

$$\text{b) } \int_0^{\pi/2} \int_0^a \frac{r}{1+r^2} dr d\theta = \frac{\pi}{2} \cdot \frac{1}{2} \ln(1+r^2) \Big|_0^a = \frac{\pi}{4} \ln(1+a^2).$$

$$\text{c) } \int_0^{\pi/4} \int_0^{\sec \theta} \tan^2 \theta \cdot r dr d\theta = \frac{1}{2} \int_0^{\pi/4} \tan^2 \theta \sec^2 \theta d\theta = \frac{1}{6} \tan^3 \theta \Big|_0^{\pi/4} = \frac{1}{6}.$$

$$\text{d) } \int_0^{\pi/2} \int_0^{\sin \theta} \frac{r}{\sqrt{1-r^2}} dr d\theta$$

$$\text{Inner: } -\sqrt{1-r^2} \Big|_0^{\sin \theta} = 1 - \cos \theta \quad \text{Outer: } \theta - \sin \theta \Big|_0^{\pi/2} = \pi/2 - 1.$$


**3B-3** a) the hemisphere is the graph of  $z = \sqrt{a^2 - x^2 - y^2} = \sqrt{a^2 - r^2}$ , so we get

$$\iint_R \sqrt{a^2 - r^2} dA = \int_0^{2\pi} \int_0^a \sqrt{a^2 - r^2} r dr d\theta = 2\pi \cdot -\frac{1}{3} (a^2 - r^2)^{3/2} \Big|_0^a = 2\pi \cdot \frac{1}{3} a^3 = \frac{2}{3} \pi a^3.$$


$$b) \int_0^{\pi/2} \int_0^a (r \cos \theta)(r \sin \theta)r \, dr \, d\theta = \int_0^a r^3 \, dr \int_0^{\pi/2} \sin \theta \cos \theta \, d\theta = \frac{a^4}{4} \cdot \frac{1}{2} = \frac{a^4}{8}.$$

c) In order to be able to use the integral formulas at the beginning of 3B, we use symmetry about the  $y$ -axis to compute the volume of just the right side, and double the answer.

$$\begin{aligned} \iint_R \sqrt{x^2 + y^2} \, dA &= 2 \int_0^{\pi/2} \int_0^{2 \sin \theta} r \, r \, dr \, d\theta = 2 \int_0^{\pi/2} \frac{1}{3} (2 \sin \theta)^3 \, d\theta \\ &= 2 \cdot \frac{8}{3} \cdot \frac{2}{3} = \frac{32}{9}, \text{ by the integral formula at the beginning of 3B.} \end{aligned}$$



$$d) 2 \int_0^{\pi/2} \int_0^{\sqrt{\cos \theta}} r^2 \, r \, dr \, d\theta = 2 \int_0^{\pi/2} \frac{1}{4} \cos^2 \theta \, d\theta = 2 \cdot \frac{1}{4} \cdot \frac{\pi}{4} = \frac{\pi}{8}.$$



### 3C. Applications of Double Integration


**3C-1** Placing the figure so its legs are on the positive  $x$ - and  $y$ -axes,

$$a) \text{ M.I.} = \int_0^a \int_0^{a-x} x^2 \, dy \, dx \quad \text{Inner: } x^2 y \Big|_0^{a-x} = x^2(a-x); \quad \text{Outer: } \left[ \frac{1}{3} x^3 a - \frac{1}{4} x^4 \right]_0^a = \frac{1}{12} a^4.$$

$$b) \iint_R (x^2 + y^2) \, dA = \iint_R x^2 \, dA + \iint_R y^2 \, dA = \frac{1}{12} a^4 + \frac{1}{12} a^4 = \frac{1}{6} a^4.$$

c) Divide the triangle symmetrically into two smaller triangles, their legs are  $\frac{a}{\sqrt{2}}$ ;

$$\text{Using the result of part (a), M.I. of } R \text{ about hypotenuse} = 2 \cdot \frac{1}{12} \left( \frac{a}{\sqrt{2}} \right)^4 = \frac{a^4}{24}$$



**3C-2** In both cases,  $\bar{x}$  is clear by symmetry; we only need  $\bar{y}$ .

$$a) \text{ Mass is } \iint_R dA = \int_0^{\pi} \sin x \, dx = 2$$

$$y\text{-moment is } \iint_R y \, dA = \int_0^{\pi} \int_0^{\sin x} y \, dy \, dx = \frac{1}{2} \int_0^{\pi} \sin^2 x \, dx = \frac{\pi}{4}; \text{ therefore } \bar{y} = \frac{\pi}{8}.$$

b) Mass is  $\iint_R y \, dA = \frac{\pi}{4}$ , by part (a). Using the formulas at the beginning of 3B,

$$y\text{-moment is } \iint_R y^2 \, dA = \int_0^{\pi} \int_0^{\sin x} y^2 \, dy \, dx = 2 \int_0^{\pi/2} \frac{\sin^3 x}{3} \, dx = 2 \cdot \frac{1}{3} \cdot \frac{2}{3} = \frac{4}{9},$$

$$\text{Therefore } \bar{y} = \frac{4}{9} \cdot \frac{4}{\pi} = \frac{16}{9\pi}.$$

**3C-3** Place the segment either horizontally or vertically, so the diameter is respectively on the  $x$  or  $y$  axis. Find the moment of half the segment and double the answer.


(a) (Horizontally, using rectangular coordinates) Note that  $a^2 - c^2 = b^2$ .

$$\int_0^b \int_c^{\sqrt{a^2-x^2}} y \, dy \, dx = \int_0^b \frac{1}{2}(a^2 - x^2 - c^2) \, dx = \frac{1}{2} \left[ b^2 x - \frac{x^3}{3} \right]_0^b = \frac{1}{3} b^3; \quad \text{ans: } \frac{2}{3} b^3.$$

(b) (Vertically, using polar coordinates). Note that  $x = c$  becomes  $r = c \sec \theta$ .

$$\text{Moment} = \int_0^\alpha \int_{c \sec \theta}^a (r \cos \theta) r \, dr \, d\theta \quad \text{Inner: } \frac{1}{3} r^3 \cos \theta \Big|_{c \sec \theta}^a = \frac{1}{3} (a^3 \cos \theta - c^3 \sec^2 \theta)$$

$$\text{Outer: } \frac{1}{3} \left[ a^3 \sin \theta - c^3 \tan \theta \right]_0^\alpha = \frac{1}{3} (a^2 b - c^2 b) = \frac{1}{3} b^3; \quad \text{ans: } \frac{2}{3} b^3.$$




**3C-4** Place the sector so its vertex is at the origin and its axis of symmetry lies along the positive  $x$ -axis. By symmetry, the center of mass lies on the  $x$ -axis, so we only need find  $\bar{x}$ .

Since  $\delta = 1$ , the area and mass of the disc are the same:  $\pi a^2 \cdot \frac{2\alpha}{2\pi} = a^2 \alpha$ .

$$x\text{-moment: } 2 \int_0^\alpha \int_0^a r \cos \theta \cdot r \, dr \, d\theta \quad \text{Inner: } \frac{2}{3} r^3 \cos \theta \Big|_0^a;$$


$$\text{Outer: } \frac{2}{3} a^3 \sin \theta \Big|_0^\alpha = \frac{2}{3} a^3 \sin \alpha \quad \bar{x} = \frac{\frac{2}{3} a^3 \sin \alpha}{a^2 \alpha} = \frac{2}{3} \cdot a \cdot \frac{\sin \alpha}{\alpha}.$$



**3C-5** By symmetry, we use just the upper half of the loop and double the answer. The upper half lies between  $\theta = 0$  and  $\theta = \pi/4$ .

$$2 \int_0^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r^2 r \, dr \, d\theta = 2 \int_0^{\pi/4} \frac{1}{4} a^4 \cos^2 2\theta \, d\theta$$

$$\text{Putting } u = 2\theta, \text{ the above} = \frac{a^4}{2 \cdot 2} \int_0^{\pi/2} \cos^2 u \, du = \frac{a^4}{4} \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \frac{\pi a^4}{16}.$$




### 3D. Changing Variables

**3D-1** Let  $u = x - 3y$ ,  $v = 2x + y$ ;  $\frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} 1 & -3 \\ 2 & 1 \end{vmatrix} = 7$ ;  $\frac{\partial(x, y)}{\partial(u, v)} = \frac{1}{7}$ .

$$\iint_R \frac{x-3y}{2x+y} \, dx \, dy = \frac{1}{7} \int_0^7 \int_1^4 \frac{u}{v} \, dv \, du$$

$$\text{Inner: } u \ln v \Big|_1^4 = u \ln 4; \quad \text{Outer: } \frac{1}{2} u^2 \ln 4 \Big|_0^7 = \frac{49 \ln 4}{2}; \quad \text{Ans: } \frac{1}{7} \frac{49 \ln 4}{2} = 7 \ln 2$$




**3D-2** Let  $u = x + y$ ,  $v = x - y$ . Then  $\frac{\partial(u, v)}{\partial(x, y)} = 2$ ;  $\frac{\partial(x, y)}{\partial(u, v)} = \frac{1}{2}$ .

To get the  $uv$ -equation of the bottom of the triangular region:

$$y = 0 \Rightarrow u = x, v = x \Rightarrow u = v.$$

$$\iint_R \cos\left(\frac{x-y}{x+y}\right) dx dy = \frac{1}{2} \int_0^2 \int_0^u \cos \frac{v}{u} dv du$$

Inner:  $u \sin \frac{v}{u} \Big|_0^u = u \sin 1$     Outer:  $\frac{1}{2} u^2 \sin 1 \Big|_0^2 = 2 \sin 1$     Ans:  $\sin 1$



**3D-3** Let  $u = x$ ,  $v = 2y$ ;  $\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{vmatrix} = \frac{1}{2}$

Letting  $R$  be the elliptical region whose boundary is  $x^2 + 4y^2 = 16$  in  $xy$ -coordinates, and  $u^2 + v^2 = 16$  in  $uv$ -coordinates (a circular disc), we have

$$\begin{aligned} \iint_R (16 - x^2 - 4y^2) dy dx &= \frac{1}{2} \iint_R (16 - u^2 - v^2) dv du \\ &= \frac{1}{2} \int_0^{2\pi} \int_0^4 (16 - r^2) r dr d\theta = \pi \left( 16 \frac{r^2}{2} - \frac{r^4}{4} \right)_0^4 = 64\pi. \end{aligned}$$

**3D-4** Let  $u = x + y$ ,  $v = 2x - 3y$ ; then  $\frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} 1 & 1 \\ 2 & -3 \end{vmatrix} = -5$ ;  $\frac{\partial(x, y)}{\partial(u, v)} = \frac{1}{5}$ .

We next express the boundary of the region  $R$  in  $uv$ -coordinates.


For the  $x$ -axis, we have  $y = 0$ , so  $u = x$ ,  $v = 2x$ , giving  $v = 2u$ .

For the  $y$ -axis, we have  $x = 0$ , so  $u = y$ ,  $v = -3y$ , giving  $v = -3u$ .

It is best to integrate first over the lines shown,  $v = c$ ; this means  $v$  is held constant, i.e., we are integrating first with respect to  $u$ . This gives

$$\iint_R (2x - 3y)^2 (x + y)^2 dx dy = \int_0^4 \int_{-v/3}^{v/2} v^2 u^2 \frac{du dv}{5}.$$

Inner:  $\frac{v^2}{15} u^3 \Big|_{-v/3}^{v/2} = \frac{v^2}{15} v^3 \left( \frac{1}{8} - \frac{-1}{27} \right)$     Outer:  $\frac{v^6}{6 \cdot 15} \left( \frac{1}{8} + \frac{1}{27} \right)_0^4 = \frac{4^6}{6 \cdot 15} \left( \frac{1}{8} + \frac{1}{27} \right)$ .




**3D-5** Let  $u = xy$ ,  $v = y/x$ ; in the other direction this gives  $y^2 = uv$ ,  $x^2 = u/v$ .

We have  $\frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} y & x \\ -y/x^2 & 1/x \end{vmatrix} = \frac{2y}{x} = 2v$ ;  $\frac{\partial(x, y)}{\partial(u, v)} = \frac{1}{2v}$ ; this gives

$$\iint_R (x^2 + y^2) dx dy = \int_0^3 \int_1^2 \left( \frac{u}{v} + uv \right) \frac{1}{2v} dv du.$$

Inner:  $\frac{-u}{2v} + \frac{u}{2} v \Big|_1^2 = u \left( -\frac{1}{4} + 1 + \frac{1}{2} - \frac{1}{2} \right) = \frac{3u}{4}$ ;    Outer:  $\frac{3}{8} u^2 \Big|_0^3 = \frac{27}{8}$ .



**3D-8** a)  $y = x^2$ ; therefore  $u = x^3$ ,  $v = x$ , which gives  $u = v^3$ .

b) We get  $\frac{u}{v} + uv = 1$ , or  $u = \frac{v}{v^2 + 1}$ ; (cf. 3D-5)